

AD-A200 304

DTIC FILE COPY

2

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0470-P00003

R&T Code NR 33359-718

Technical Report No. 100

He(I) Photoelectron Spectrum of Methylcyclopropene Derivatives

by

T. Koenig, T. Curtess, R. Winter, S. Pons, K. Ashley, Q. Mei, P. Stang,
B. Halton, S.J. Buckland and D. R. Rolison

Prepared for publication in J. Org. Chem.

Department of Chemistry
University of Utah
Salt Lake City, UT 84112

July 15, 1988

DTIC
ELECTE
NOV 14 1988
S D
CH

Reproduction in whole, or in part, is permitted for
any purpose of the United States Government

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

88 11 10 062

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release and sale. Distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) ONR Technical Report No. 100		7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION University of Utah	6b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code)	
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Henry Eyring Building Salt Lake City, UT 84112		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-83-K-0470-P00003	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (if applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program, Code 1113 800 N. Quincy Street Arlington, VA 22217		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) He(I) Photoelectron Spectrum of Methylcyclopropane Derivatives			
12. PERSONAL AUTHOR(S) T. Koenig, T. Curless, R. Winter, S. Pons, K. Ashley, Q. Mei, P. Stang, B. Halton, J. Buckland and D. R. Rolison			
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM 9/87 TO 7/88	14. DATE OF REPORT (Year, Month, Day) July 15, 1988	15. PAGE COUNT 24
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Attached.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Stanley Pons		22b. TELEPHONE (Include Area Code) (801)581-4760	22c. OFFICE SYMBOL

Abstract

The He(I) photoelectron spectra of benzo and naphtho derivatives of 1,1-diphenylmethylenecyclopropene (DPMC) are presented. A correlation of electrochemical oxidation potentials (E_{sub}^{+} values) and gas phase ionization potentials of a series of aromatic hydrocarbons is discussed. This correlation suggests that the difference in E_{sub}^{+} values for the two DPMC derivatives is due to structure specific solvation of the radical cations. HAM/3 and HAM/3/CI calculations are compared with a simple structure derived analysis of the observed pe spectra. It is suggested that the effects of the fused cyclopropene grouping are to be found in small (~ 2 eV) reductions of the second or higher ionization energies. (A.D.)_R



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or
A-1	

THE HE(I) PHOTOELECTRON SPECTRA OF METHYLENECYCLOPROPENE DERIVATIVES
CORRELATION WITH ELECTROCHEMICAL OXIDATION

T. Koenig*, Tim Curtiss and Rolf Winter

Department of Chemistry
University of Oregon
Eugene, Oregon 97403
U.S.A.

Kevin Ashley, Qiu Mei, Peter J. Stang and Stanley Pons


Department of Chemistry
University of Utah
Salt Lake City, Utah 84112
U.S.A.

Simon J. Buckland and Brian Halton

Department of Chemistry
Victoria University of Wellington
Wellington, New Zealand

Introduction

Methylenecyclopropene (**1**), the smallest non-alternant hydrocarbon, has recently been prepared¹ and spectral observations² have confirmed the existence of a significant dipolar (**1a**) contribution to its ground state electronic structure. Characterization of $\mathbf{1} \longleftrightarrow \mathbf{1a}$ is complicated by its moderately



high chemical reactivity but relatively complete spectral and theoretical determinations for this system are possible and forthcoming.³ The present work concerns results on the kinetically stable cycloproparenes **2** and **3**, the diarylated benzo and naphtho derivatives of **1**.⁴



The stability, imparted by the aryl substitution, allowed some of us to study⁵ the electrochemical behavior of **2** and **3**. A significant result of that work was the fact that the half wave oxidation potential (E_h^{+}) for **3** (0.81 eV) was higher than that for **2** (0.68 eV). This ordering goes against the usual idea that the more delocalized system (**3**) should be more easily oxidized. It thus seemed of interest to investigate the photoelectron spectra (pe) of **2** and **3** to see if the E_h^{+} difference corresponded to an ionization potential (IP) difference in the gas phase. We now report on our determination of the pe spectra of **2** and **3** and on an analysis of the first several ionic states for these systems. A general correlation between E_h^{+} values and

pes IP's for aromatic hydrocarbons is presented. This correlation suggests that solvation effects, in the electrochemical oxidation of arenes, increase with increasing IP, due to the smaller molecular size that attends the increased IP with in the series. The difference in E_{H}^+ values for 2 and 3 is rationalized as residual differential solvation effects arising from the difference in hole density distribution.

Experimental

The syntheses of 2 and 3 have been published.⁴ The half wave oxidation potentials for 2 and 3 were also as previously described.⁵ The photoelectron spectra were obtained from our ⁶ $\pi/2$ sector instrument at temperatures between 50 and 100°C. The spectra were calibrated in duplicate versus Ar, Xe, methyl iodide mixtures. The beginning resolution in each pe determination was less than 30 mev.

Results and Discussion

The pe spectra of **2** and **3** are shown in Figure 1. They indicate that the observed difference in E_h^+ values has no clear counterpart in the gas phase. The first ionization potential of **2** is essentially the same as **3**. This applies to both the first peak maximum ($-IP$ (vertical), $7.145 \pm .03$ eV) and onset ($-IP$ adiabatic, $6.84 \pm .03$ eV). Figure 2 shows a plot of E_h^+ values^{5,7a} against first IP values^{7,8} (vertical) for several aromatic systems. The correlation predicts an E_h^+ value of 0.74 eV for both **2** and **3**. The observed value for **2** is slightly lower (0.68 eV) while that for **3** is slightly higher (0.81 eV) than the value (0.74 eV) that the linear correlation predicts.




Figure 2. Correlation of E_h^+ with IP.

The regression line (correlation coefficient 0.989) shown in Figure 2 is expressed by equation (1). This can be compared with the simplistic theor-

$$E_h^+ = 0.667 \cdot IP - 4.023 \quad (1)$$

etical expression given in equation (2) where $\Delta\Delta G_{sol}^{\circ}$ is the difference in

$$E_h^+ = IP + \Delta\Delta G_{sol}^{\circ} + \text{constant} \quad (2)$$

solvation of the neutral (reduced) and ionic (oxidized) forms of the hydrocarbon and the constant corrects for the reference electrode potential. The linearity and less than unit slope in equation (1) suggest that $\Delta\Delta G_{sol}^{\circ}$ contains a term that is proportional to IP. Equation 3 incorporates this observation and defines the residual differential solvation, $\delta\Delta\Delta G_{sol}^{\circ}$.

$$\Delta\Delta G_{sol}^{\circ} = -S \cdot IP + \delta\Delta\Delta G_{sol}^{\circ} \quad (3)$$

Equation (4) results from these definitions.


$$E_h^+ = (1-S) \cdot IP + \delta\Delta\Delta G_{sol}^{\circ} + \text{constant} \quad (4)$$

The empirical relationship (equation 1) is obtained by neglecting the $\delta\Delta\Delta G_{sol}^{\circ}$ term with s equal to 0.333 and the constant equal to -4.023 (eV). The deviations from the value (0.74 eV) predicted for 2 and 3 can be interpreted as residual differential solvation effects ($\delta\Delta\Delta G_{sol}^{\circ}$). The values of these deviations (-0.06 eV for 2 and +0.07 eV for 3) are not atypical of the other members of the series shown in Figure 2. Small structure specific solvation effects are certainly present and the proposed $\delta\Delta\Delta G_{sol}^{\circ}$ values are one way to recognize them at least phenomenologically.

The $-S \cdot IP$ term of equation (4) may be related to recent observations⁹ of the E_h^+ values of a series of molecules of high IP. In those studies it was found that the correlation between E_h^+ and IP changed from one which was linear and of unit slope (saturated hydrocarbons) to one of less than unit slope at the high IP -- small molecular size end. The assistance to oxidation by the solvent is of greater magnitude in the high IP - small molecular size

range. Reduced molecular size accompanies increased IP in the aromatic hydrocarbon series of Figure 2 and the $-S \cdot IP$ term of equation (4) may be explained by the higher vacancy density per unit volume in the ions of high IP and smaller π orbital size. The difference in E_N^+ for 2 and 3 are well within expectations for structure specific solvation energy differences which we describe by the $\delta\Delta G_{sol}^+$ term.

SCHEME I



The observed (Figure 1) near equality in the first IP's of 2 and 3 can be understood by considering their π electron systems as perturbed¹⁰ 1,1-diphenylethenes.¹¹ The energetic positions and symmetries of the perturbing benzyne (2) and naphthynic (3) π electron groups, with respect to that of the lowest ionic state of 1,1-diphenylethene¹¹ (4), are shown in Scheme I. The conrotation of the two phenyl groups with respect to coplanarity, shown by the X-ray structure for 2, leaves an approximate¹² C_2 axis along the exo methylene group. The labelling in Scheme I designates the basis structures as symmetric (S_B , 2; S_N , 3) or antisymmetric (A_B , 2, A_N , 3; A_E , 4) with respect to this symmetry element. The first IP's of 2 and 3 are then describable as combinations from the appropriate A_1 manifold.




The first naphtho (A_u , Scheme I) hole structure in 3, that will perturb the first one derived from 4 (A_g , 8.25 eV,¹¹ Scheme I), corresponds to the second ionic state of naphthalene ($^2B_{1u}$, 8.88 eV⁸). The corresponding hole structure for 2 (A_g , Scheme I) is one of the e_g pair of benzene (9.3 eV). The difference in energy between A_u and A_g is 0.6 eV (3) while that for A_g and A_g is 1.02 eV (2). The smaller difference for 3 compared to 2 leads one to expect a lower first IP for 3 as is the usual case. The interaction constant (H_{AA} , Scheme I) for mixing A_u and A_g (3) will be smaller than that for A_g and A_g (2). The lowered interaction constant for 3 would lead to a high first IP in 3 compared to 2. The net result of the smaller energy gap and smaller interaction constant for 3 is the fortuitous equality in the first IP of 3 with that of 2.

In this very simple analysis, the ionic ground states of 2 and 3 would both be completely (π) delocalized. However, the fortuitous equality in first IP's of 2 and 3 implies that essentially the same (~70%) fraction of the vacancy should reside on the 1,1-diphenylethene grouping in the two ionic ground states. The fraction of the hole distributed on the benzo and naphtho grouping would also be the same (30%). The vacancy density, on a per atom basis, would be lower in 3 compared to 2 because of the large size of the naphtho substructure. This would be expected to lead to a lowered differential solvation energy ($\delta\Delta G_{sol}^*$) and hence an increased E_u^* for 3 compared to 2. This explanation of the difference in the E_u^* values of 2 compared to 3 is at least internally consistent since it is derived from the same physical source as suggested for the larger -S·IP term in equation (4).

Scheme I also predicts that the second pe band for 3 should correspond to the first ionic state of naphthalene (8.15 eV, $^2A_u^8$). The symmetry of this basis representation (S_u) is such that it will not be overlap-sensitive to the

attached 1,1-diphenylethene unit. The second ionic state of 3 should thus be describable as one localized on the naphtho unit with bond distance changes that correspond to those for the lowest ionic state of naphthalene itself. Indeed, the position (7.95 eV) and vibrational shape of the second pe band of 3 (Figure 1) strongly support this assignment.

Analysis of the pe spectra of 2 and 3 in terms of perturbed diphenylethene (4) ionic states can also be extended to the 9 eV region. The experimental spectrum of 4¹¹ shows that two ionic states are located (both) at 9.05 eV with a third state at 9.2 eV. This set is describable in terms of combinations of hole structures localized on the two phenyl groups with little or no ethene (π) cationic character. Two of the three are of S symmetry (S_{ϕ_2} (1), S_{ϕ_2} (2) below). The one of A symmetry (A_{ϕ_2}) has low coefficients at the phenyl carbons that are joined to the ethene group giving negligible ethene-phenyl mixing.



The S_{ϕ_2} structure (9.3 eV) for 2, shown in Scheme I, is likewise isolated by symmetry from the ethene unit. Thus, for 2, at least 4 ionic states should reside in the 9 eV region. The observed intensity pattern in the spectrum of 2 supports this analysis since the relative area of the 8.7-9.9 eV section is 4.5 times that of the resolved 7.14 eV band. These arguments also apply to 3 (except for the S_{ϕ_2} structure) and the gross appearance of the 9 eV region of the pe spectrum of 3 is expected to be similar to that for 2.

While the major features of the low energy region of the pe spectra of 2 and 3 are simply understandable, the above analysis has ignored possible

effects due to dipolar contributions analogous to 1a in the neutral hydrocarbons and effects due to distortions of the fused benzo and naphtho groups. The X-ray data for 2⁴ show a significant reduction in the bond distance between the two carbons in the fused cyclopropene ring. Table 1 shows results of HAM/3¹³ and HAM/3/CI¹⁴ calculations for 2 and 3. The X-ray structure was used for 2 and the additional C-H groups of 3 were simply added at naphthalene positions. These calculations predict essentially equal first ionization potentials for 2 and 3 although the calculated values are appreciably higher than observed. The high values are not too surprising since the HAM/3 transition state method overestimates the first IP of benzene by a similar (~0.4 eV) amount. The HAM/3 method has not been previously tested on such large systems and the results of Table 1 suggest that the existing parameterization may not yet be optimal for such systems.

The position of the S_u state of 3 (7.95 eV) is shifted downwards compared to the corresponding band in the spectrum of naphthalene (8.15 eV).⁸ This decrease is in the direction expected for the shortened (1.355Å) C-C distance between the two carbon atoms common to the cyclopropene and naphthalene rings. The HAM/3 calculations give a calculated shift of -0.19 eV in the S_u band for 3 compared to naphthalene at its experimental geometry. A similar shift is calculated for the S_g position in 2 compared to benzene but the S_g band is not resolved from the other 4 states seen in the 8.75-9.7 eV group for 2. The position of the S_u band in the spectrum of 3 may be taken as a measure of the net shift attributable to the shortened bond length in the ring fusing carbon atoms in 3.

The direction of the dipole (1a) of 1 suggests that the ionic states that are localized on the two phenyl substituents in 2 and 3 might be stabilized so that they would appear at lower IP in 2 and 3 than in 4. The observed spectra

could be viewed as being in agreement with this expectation in that the maxima in the second π bands of 2 (8.7 eV) and the third band in 3 (8.8 eV) are lower than that corresponding in 4 (9.05 eV) by 0.2-0.3 eV. The HAM/3 calculated shifts for these (S_{ϕ_2} , A_{ϕ_2} , Table 1) states, in 2 and 3 compared to 4, are between -0.24 eV and -0.34 eV. The small reduction in the positions of the second and third bands in the spectra of 2 and 3 could thus be a reflection of the increased polarization or polarizability of the ethene group in 2 and 3 compared to 4 and thus reflect the nonalternacy of the system.

The calculations at detailed geometries confirm the simple analysis given above in terms of band assignments up to the -9.5 eV point of the observed spectra. However, the simple analysis ignored some of the basis functions in the A_1 manifold which are included in the HAM/3 calculations. These include the basis function associated with the 10.26 eV band in the spectrum of 4¹¹ which we label A'_1 in Table 2. Also, the naphtho compound (3) should show a second symmetry factored (S) state which corresponds to the third (${}^2B_{2g}$) ionic state in naphthalene (10.08 eV). This state for 3 is labeled S'_M in Table 1. The calculated position of the first member of the σ manifold is also included in Table 1. The HAM/3 calculations and assignments seem to be in acceptable agreement with the experimental spectra.

The HAM/3/CI¹⁴ results (Table 1) suggest that non-Koopmans' effects are appreciable in the ionic states of 2 and 3 beginning in the 10-11 eV region. Such shake-up phenomena give an increased number of bands which are broader than single excitation counterparts. The observed spectra of 2 and 3, between 10 and 11 eV, mildly suggest that these effects may be present but the number of states is too high to show shake up effects very clearly. As has been discussed¹⁵ in the case of stilbene, the rotational angle of the phenyl groups with respect to the ethene group in 2, 3 and 4 may not be the same as given by

crystal structures. Additional calculations on these relative large systems at variable geometries would be required before their pe spectra can be analyzed to the non-Koopmans (configuration interaction) level. Other derivatives such as the fluorenylidene analogs of 2 and 3, would be better candidates for revealing the non-Koopmans effects.

Conclusions

The present work shows that there is little or no difference in the first ionization potentials of 2 and 3. An apparently linear correlation between E_h^+ values and gas phase IP's of a series of aromatic hydrocarbons indicates that the difference in E_h^+ values of 2 and 3 is due to structure specific solvation effects. The HAM/3 method gives acceptable agreement with the experimental spectra in terms of order of states but appears to overestimate all of the IP's to a fairly significant extent. The HAM/3/CI method indicates that shakeup phenomena are expected in this series above the 9.5 eV level. Studies of other members of this series could serve to test the E_h^+ -IP correlation and might show the non-Koopmans effects more clearly.

Acknowledgement


S. P. and K. A. thank the Office of Naval Research for support of this work. P. S. and T. K. thank the NSF (CHE 84-19099, CHE 84-19950) for their support.

References


1. Staley, S. W.; Norden, T. D. J. Am. Chem. Soc. 1984, 106, 3699; Billups, W. E.; Lin, L. J.; Casserly, E. W. ibid. 1984, 106, 3698; Maier, G.; Hoppe, M.; Lanz, K.; Reisenauer, P. Tetrahedron Lett. 1984, 25, 5465.
2. Norden, Timothy D.; Staley, Stuart W.; Taylor, William H.; Harmony, Marlin D. J. Am. Chem. Soc. 1986, 108, 7912-7918.
3. An account of the observed photoelectron spectrum of 1 is currently submitted. S. Staley, private communication.
4. Halton, Brian; Randall, Clifford J.; Gainsford, Graeme J.; Stang, Peter J. J. Am. Chem. Soc. 1986, 108, 5949-5956.
5. Ashley, Kevin; Foley, John K.; Mei, Qui; Ghoroghchian, Jamal; Sarfarazi, Fereshteh; Cassidy, John; Halton, Brian; Stang, Peter; Pons, Stanely J. Org. Chem. 1986, 51, 2089; Ashley, K.; Sarfarazi, F.; Buckland, S. J.; Foley, J. K.; Mei, Q.; Halton, B.; Stang, P. J.; Pons, S. Can. J. Chem., in press.
6. Imre, D. and Koenig, T. Chem. Phys. Lett. 1980, 23, 62.
7. Koenig, T.; Tuttle, M. and Wielesek, R. A. Tetrahedron Lett. 1974, 2537-2540; Koenig, T. and Tuttle, M. J. Org. Chem. 1974, 39, 1308.
- 7a. Ross, S. D.; Finkelstein, M.; Rudd, E. F., "Anodic Oxidation", in Organic Chemistry. A Series of Monographs, Vol. 32, Academic Press: New York, 1975, Ch. 5; Ebersson, E.; Utley, J. H. P., in "Organic Electrochemistry", 2nd ed. M. M. Baiser and H. Lund, eds., Marcel Dekker: New York, Ch. 13; Yoshida, K., "Electrooxidation in Organic Chemistry. The Role of Cation Radicals as Synthetic Intermediates," Wiley: New York, 1984; Phelps, J.; Santhanam, K. S. V.; Bard, A. J. J. Am. Chem. Soc. 1967, 81, 1752.
8. Heilbronner, Edgar; Maier, J. P., in "Electron Spectroscopy. Theory, Techniques and Applications," Brundle, C. R.; Baker, A. D., eds., Academic, New York, 1977, Vol. 1, pp 205-287.
9. Dibble, T.; Bandyopadhyay, S.; Ghoroghchian, J.; Smith, J. J.; Sarfarazi, F.; Fleischmann, M.; Pons, J. J. Phys. Chem. 1986, 90, 5275.
10. Koenig, T.; Imre, Daniel; Hoobler, James A. J. Am. Chem. Soc. 1979, 101, 6446; Koenig, T. and Longmaid, H. J. Org. Chem. 1974, 39, 560.
11. Maier, J. P. and Turner, D. W. J. Chem. Soc. Faraday II, 1973, 69, 196-206.
12. The X-ray structure of 2 indicates non-planarity and distortion of the cyclopropene ring. In spite of the absence of any real symmetry elements, the approximate C₂ axis along the ethene bond is useful.
13. Lindholm, L.; Asbrink, L., "Molecular Orbitals and Their Energies by the Semiempirical HAM Method," Springer Verlag, Berlin, 1985.

14. Koenig, T.; Winter, R. and Rudolf, K. J. Am. Chem. Soc. 1987, 109, 2515.
15. Bigelow, Richard W. Chem. Phys. Lett. 1985, 117 (1), 22-8.

TABLE 1. Calculated and Observed Outer Valence PE Spectra of 2 and 3.



2



3

Γ^a	obs (eV)	HAM/3	HAM3/CI ^b	obs	HAM/3	HAM3/CI ^b
A	7.15	7.67	7.53	7.14	7.66	7.59
S _N				7.95	8.21	8.14
S _{$\phi_2(1)$} ^c	8.75	8.97	9.02	8.91	8.88	9.09
A _{ϕ_2} ^c	d	9.00	9.10	d	8.92	9.17
S _{$\phi_2(2)$} ^c	9.12	9.08	9.11	9.08	9.00	9.19
S _B	d	9.52	9.83			
A _B or A _N	d	9.67	9.79	d	9.40	9.33
A _E '	10.2	10.64	{ 10.80* 11.06*	9.95	10.02	{ 9.99* 10.55* 10.67*
S _N '				d	10.05	10.37
σ	10.8	10.79		(10.7)	10.67	

^aApproximate symmetry. ^bA * designates a state with appreciable shakeup character (Ref. 14). ^cState with vacancy localized on the diphenyl substituent. ^dPeak unresolved but present.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

ABSTRACTS DISTRIBUTION LIST, SDIO/IST

Dr. Robert A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, NY 14214

Dr. Douglas N. Bennion
Department of Chemical Engineering
Brigham Young University
Provo, UT 84602

Dr. Stanley Pons
Department of Chemistry
University of Utah
Salt Lake City, UT 84112

Dr. H. V. Venkatesetty
Honeywell, Inc.
10701 Lyndale Avenue South
Bloomington, MN 55420

Dr. J. Foos
EIC Labs Inc.
111 Downey St.
Norwood, MA 02062

Dr. Neill Weber
Ceramatec, Inc.
163 West 1700 South
Salt Lake City, UT 84115

Dr. Subhash C. Narang
SRI International
333 Ravenswood Ave.
Menlo Park, CA 94025

Dr. J. Paul Pemsler
Castle Technology Corporation
52 Dragon Ct.
Woburn, MA 01801

Dr. R. David Rauh
EIC Laboratory Inc.
111 Downey Street
Norwood, MA 02062

Dr. Joseph S. Foos
EIC Laboratories, Inc.
111 Downey Street
Norwood, Massachusetts 02062

Dr. Donald M. Schleich
Department of Chemistry
Polytechnic Institute of New York
333 Jay Street
Brooklyn, New York 01

Dr. Stan Szpak
Code 633
Naval Ocean Systems Center
San Diego, CA 92152-5000

Dr. George Blomgren
Battery Products Division
Union Carbide Corporation
25225 Detroit Rd.
Westlake, OH 44145

Dr. Ernest Yeager
Case Center for Electrochemical
Science
Case Western Reserve University
Cleveland, OH 44106

Dr. Mel Miles
Code 3852
Naval Weapons Center
China Lake, CA 93555

Dr. Ashok V. Joshi
Ceramatec, Inc.
2425 South 900 West
Salt Lake City, Utah 84119

Dr. W. Anderson
Department of Electrical &
Computer Engineering
SUNY - Buffalo
Amherst, Massachusetts 14260

Dr. M. L. Gopikanth
Chemtech Systems, Inc.
P.O. Box 1067
Burlington, MA 01803

Dr. H. F. Gibbard
Power Conversion, Inc.
495 Boulevard
Elmwood Park, New Jersey 07407

ABSTRACTS DISTRIBUTION LIST, SDIO/IST

Dr. V. R. Koch
Covalent Associates
52 Dragon Court
Woburn, MA 01801

Dr. Randall B. Olsen
Chronos Research Laboratories, Inc.
4186 Sorrento Valley Blvd.
Suite H
San Diego, CA 92121

Dr. Alan Hooper
Applied Electrochemistry Centre
Harwell Laboratory
Oxfordshire, OX11 0RA UK

Dr. John S. Wilkes
Department of the Air Force
The Frank J. Seiler Research Lab.
United States Air Force Academy
Colorado Springs, CO 80840-6528

Dr. Gary Bullard
Pinnacle Research Institute, Inc.
10432 N. Tantan Avenue
Cupertino, CA 95014

Dr. J. O'M. Bockris
Ementech, Inc.
Route 5, Box 946
College Station, TX 77840

Dr. Michael Binder
Electrochemical Research Branch
Power Sources Division
U.S. Army Laboratory Command
Fort Monmouth, New Jersey 07703-5000

Professor Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton, Hants, SO9 5NH UK